

PR-4

AN EFFICIENT SYNTHESIS OF POLYCYCLIC AROMATIC HYDROCARBON-CONTAINING COPOLYMERS


A. Mukherjee,^{1*} I. S. Kovalev,^{1,2} D. S. Kopchuk,^{1,2} S. Santra,¹ G. V. Zyryanov,^{1,2} O. N. Chupakhin^{1,2}

¹*Ural Federal University, 19 Mira St, Yekaterinburg, 620002, Russia.*

²*Ya. Postovsky Institute of Organic Synthesis UB RAS,
20/22 S. Kovalevskoy / Akademicheskaya St, Yekaterinburg, 620990, Russia.*

*E-mail: anindita0423@gmail.com

Abstract. The derivatives of polycyclic aromatic hydrocarbon (PAH) display unique properties of fluorescence like excellent photostability,¹ lifetime of long fluorescence and including high quantum yield. Polycyclic aromatic hydrocarbon (PAH) could be used for organic light emitting diodes, biological fluorescent probe, pressure-sensitive materials etc.² In the polymer materials area, here PAH is established into the polymer to label macromolecular chains, modify carbon nanotubes³ or improve polymer fluorescence as well. In this work we are glad to inform an efficient approach for the synthesis of PAH (such as pyrene, anthracene and naphthalene) containing copolymers based on acrylic acid. First, for the synthesis of starting vinyl PAH, we have done vinylation of the PAH. Therefore, with respect to the starting vinyl PAH we used various proportion of acrylic acid.



PAH = naphthalene, anthracene, pyrene

Figure 1. PAH containing synthesized polymers.

References

1. Maeda, H. Alkynylpyrenes as Improved Pyrene-Based Biomolecular Probes with the Advantages of High Fluorescence Quantum Yields and Long Absorption/Emission Wavelengths / H. Maeda, T. Maeda, K. Mizuno, K. Fujimoto, H. Shimizu, M. Inouye// *Chemistry - A European Journal*. – 2006. – Vol. 12. Iss. 3. – P. 824–831.
2. Diring, S. Luminescent Ethynyl-Pyrene Liquid Crystals and Gels for Optoelectronic Devices / S. Diring, F. Camerel, B. Donnio, T. Dintzer, S. Toffanin, R. Capelli, M. Muccini, R. Ziessel// *Journal of the American Chemical Society*. – 2009. – Vol. 131. Iss. 50. – P. 18177–18185.
3. Yuan, W. Z. Electronic Interactions and Polymer Effect in the Functionalization and Solvation of Carbon Nanotubes by Pyrene- and Ferrocene-Containing Poly(1-alkyne)s / W. Z. Yuan, Y. Mao, H. Zhao, J. Z. Sun, H. P. Xu, J. K. Jin, Q. Zheng, B. Z. Tang// *Macromolecules*. 2008. – Vol. 41. Iss. 3. – P. 701–707.

This work was supported by Russian Science Foundation (Ref no. 20-73-10205) and the Russian Foundation for Basic Research (grant # 19-53-55002)